Rawls et al. (1983) J. Hydraul. Eng. 109:62-70

TABLE 2.—Green and Ampt Parameters According to Soil Texture Classes and Horizons

Soil texture class (1)	Horizon (2)	Sample size (3)	Total porosity, ϕ, in cubic centimeters per cubic centimeters (4)	Effective porosity, θ_e, in cubic centimeters per cubic centimeters (5)	Wetted front capillary pressure, ψ_f,[a] in centimeters (6)	Hydraulic conductivity, K,[b] in centimeters per hour (7)
Sand[c]		762	0.437 0.374–0.500[d]	0.417 (0.354–0.480)	4.95 (0.97–25.36)	11.78
	A	370	0.452 (0.396–0.508)	0.431 (0.375–0.487)	5.34 (1.24–23.06)	
	B	185	0.440 (0.385–0.495)	0.421 (0.365–0.477)	6.38 (1.31–31.06)	
	C	127	0.424 (0.385–0.463)	0.408 (0.365–0.451)	2.07 (0.32–13.26)	
Loamy sand		338	0.437 (0.363–0.506)	0.401 (0.329–0.473)	6.13 (1.35–27.94)	2.99
	A	110	0.457 (0.385–0.529)	0.424 (0.347–0.501)	6.01 (1.58–22.87)	
	B	49	0.447 (0.379–0.515)	0.412 (0.334–0.490)	4.21 (1.03–17.24)	
	C	36	0.424 (0.372–0.476)	0.385 (0.323–0.447)	5.16 (0.76–34.85)	
Sandy loam		666	0.453 (0.351–0.555)	0.412 (0.283–0.541)	11.01 (2.67–45.47)	1.09
	A	119	0.505 (0.399–0.611)	0.469 (0.330–0.608)	15.24 (5.56–41.76)	
	B	219	0.466 (0.352–0.580)	0.428 (0.271–0.585)	8.89 (2.02–39.06)	
	C	66	0.418 (0.352–0.484)	0.389 (0.310–0.468)	6.79 (1.16–39.65)	
Loam		383	0.463 (0.375–0.551)	0.434 (0.334–0.534)	8.89 (1.33–59.38)	0.34
	A	76	0.512 (0.427–0.597)	0.476 (0.376–0.576)	10.01 (2.14–46.81)	
	B	67	0.512 (0.408–0.616)	0.498 (0.382–0.614)	6.40 (1.01–40.49)	
	C	47	0.412 (0.350–0.474)	0.382 (0.305–0.459)	9.27 (0.87–99.29)	
Silt loam		1,206	0.501 (0.420–0.582)	0.486 (0.394–0.578)	16.68 (2.92–95.39)	0.65
	A	361	0.527 (0.444–0.610)	0.514 (0.425–0.603)	10.91 (1.89–63.05)	
	B	267	0.533 (0.430–0.636)	0.515 (0.387–0.643)	7.21 (0.86–60.82)	
	C	73	0.470 (0.409–0.531)	0.460 (0.396–0.524)	12.62 (3.94–40.45)	
Sandy clay loam		498	0.398 (0.332–0.464)	0.330 (0.235–0.425)	21.85 (4.42–108.0)	0.15
	A	—[e]	— —	— —	— —	
	B	198	0.393 (0.310–0.476)	0.330 (0.223–0.437)	26.10 (4.79–142.30)	
	C	32	0.407 (0.359–0.455)	0.332 (0.251–0.413)	23.90 (5.51–103.75)	
Clay loam		366	0.464 (0.409–0.519)	0.309 (0.279–0.501)	20.88 (4.79–91.10)	0.10
	A	28	0.497 (0.434–0.560)	0.430 (0.328–0.532)	27.00 (6.13–118.9)	
	B	99	0.451 (0.401–0.501)	0.397 (0.228–0.530)	18.52 (4.36–78.73)	
	C	55	0.452 (0.412–0.492)	0.400 (0.320–0.480)	15.21 (3.79–61.01)	
Silty clay loam		689	0.471 (0.418–0.524)	0.432 (0.347–0.517)	27.30 (5.67–131.50)	0.10
	A	65	0.509 (0.449–0.569)	0.477 (0.410–0.544)	13.97 (4.20–46.53)	
	B	191	0.469 (0.423–0.515)	0.441 (0.374–0.508)	18.56 (4.08–84.44)	
	C	39	0.475 (0.436–0.514)	0.451 (0.386–0.516)	21.54 (4.56–101.7)	
Sandy clay		45	0.430 (0.370–0.490)	0.321 (0.207–0.435)	23.90 (4.08–140.2)	0.06
	A	—	— —	— —	— —	
	B	23	0.435 (0.371–0.499)	0.335 (0.220–0.450)	36.74 (8.33–162.1)	
	C	—	— —	— —	— —	
Silty clay		127	0.479 (0.425–0.533)	0.423 (0.334–0.512)	29.22 (6.13–139.4)	0.05
	A	—	— —	— —	— —	
	B	38	0.476 (0.445–0.507)	0.424 (0.345–0.503)	30.66 (7.15–131.5)	
	C	21	0.464 (0.430–0.498)	0.416 (0.346–0.486)	45.65 (18.27–114.1)	
Clay		291	0.475 (0.427–0.523)	0.385 (0.269–0.501)	31.63 (6.39–156.5)	0.03
	A	—	— —	— —	— —	
	B	70	0.470 (0.426–0.514)	0.412 (0.309–0.515)	27.72 (6.21–123.7)	
	C	23	0.483 (0.441–0.525)	0.419 (0.294–0.544)	54.65 (10.59–282.0)	

[a]Antilog of the log mean and standard deviation.
[b]Values for Rawls, et al. (13).
[c]Values for the texture class.
[d]Numbers in ( ) ± one standard deviation.
[e]Insufficient sample to determine parameters.